

## Product datasheet for **TP720876L**

### **CXCL5 (NM\_002994) Human Recombinant Protein**

#### **Product data:**

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human chemokine (C-X-C motif) ligand 5 (CXCL5)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Leu44-Asn114
Tag:	Tag Free
Predicted MW:	7.95 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<a href="#">NP_002985</a>
Locus ID:	6374
UniProt ID:	<a href="#">P42830</a> , <a href="#">Q6I9S7</a>
RefSeq Size:	2505
Cytogenetics:	4q13.3
RefSeq ORF:	342
Synonyms:	ENA-78; SCYB5



[View online »](#)

<b>Summary:</b>	This gene encodes a protein that is a member of the CXC subfamily of chemokines. Chemokines, which recruit and activate leukocytes, are classified by function (inflammatory or homeostatic) or by structure. This protein is proposed to bind the G-protein coupled receptor chemokine (C-X-C motif) receptor 2 to recruit neutrophils, to promote angiogenesis and to remodel connective tissues. This protein is thought to play a role in cancer cell proliferation, migration, and invasion. [provided by RefSeq, May 2013]
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Protein Pathways:</b>	Chemokine signaling pathway, Cytokine-cytokine receptor interaction