

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP720895XL

Macrophage Inflammatory Protein 4 (CCL18) (NM_002988) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated) (CCL18)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Ala21-Ala89
Tag:	N-His
Predicted MW:	10.1 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Reconstitution Method: Storage:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less
	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling
Storage: Stability:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions.
Storage: Stability: RefSeq:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 002979</u>
Storage: Stability: RefSeq: Locus ID:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 002979</u> 6362
Storage: Stability: RefSeq: Locus ID: UniProt ID:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 002979</u> 6362 <u>P55774</u>
Storage: Stability: RefSeq: Locus ID: UniProt ID: RefSeq Size:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 002979</u> 6362 <u>P55774</u> 798



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Macrophage Inflammatory Protein 4 (CCL18) (NM_002988) Human Recombinant Protein – TP720895XL
Summary:	This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for naive T cells, CD4+ and CD8+ T cells and nonactivated lymphocytes, but not for monocytes or granulocytes. This chemokine attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes. It may play a role in both humoral and cell-mediated immunity responses. [provided by RefSeq, Sep 2014]
Protein Families Protein Pathwa	

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US