

# Product datasheet for TP720898L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **UBE2C (NM 181801) Human Recombinant Protein**

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human ubiquitin-conjugating enzyme E2C (UBE2C),

transcript variant 4

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Ile141-Ser188

Tag: N-Trx&His Predicted MW: 23.3 kDa

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 3 months from date of receipt under proper storage and handling

conditions.

RefSeg: NP 861517

 Locus ID:
 11065

 UniProt ID:
 000762

 RefSeq Size:
 901

Cytogenetics: 20q13.12

RefSeq ORF: 540

**Synonyms:** dJ447F3.2; UBCH10





#### UBE2C (NM\_181801) Human Recombinant Protein - TP720898L

**Summary:** The modification of proteins with ubiquitin is an important cellular mechanism for targeting

abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]

**Protein Families:** Druggable Genome

**Protein Pathways:** Ubiquitin mediated proteolysis