

## **Product datasheet for TP720929**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Hsp40 (DNAJB1) (NM\_006145) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human DnaJ (Hsp40) homolog, subfamily B, member 1

(DNAJB1)

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Gly2-Ile340

Tag: C-His

**Predicted MW:** 39.1 kDa **Concentration:** lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

**Stability:** Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** <u>NP 006136</u>

**Locus ID:** 3337

UniProt ID: <u>P25685, Q6FHS4</u>

RefSeq Size: 2419

Cytogenetics: 19p13.12

RefSeq ORF: 1020

Synonyms: Hdj1; Hsp40; HSPF1; RSPH16B; Sis1







**Summary:** 

This gene encodes a member of the DnaJ or Hsp40 (heat shock protein 40 kD) family of proteins. DNAJ family members are characterized by a highly conserved amino acid stretch called the 'J-domain' and function as one of the two major classes of molecular chaperones involved in a wide range of cellular events, such as protein folding and oligomeric protein complex assembly. The encoded protein is a molecular chaperone that stimulates the ATPase activity of Hsp70 heat-shock proteins in order to promote protein folding and prevent misfolded protein aggregation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]