

Product datasheet for TP720967XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Neurokinin B (TAC3) (NM 013251) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human tachykinin 3 (TAC3), transcript variant 1

Species: Human
Expression Host: E. coli

Expression cDNA Clone

Gln17-Glu121

or AA Sequence:

Tag: N-His

Predicted MW: 13.9 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 037383

Locus ID: 6866

UniProt ID: Q9UHF0, A0A024RB47

RefSeq Size: 841

Cytogenetics: 12q13.3

RefSeq ORF: 363

Synonyms: HH10; LncZBTB39; NK3; NKB; NKNB; PRO1155; ZNEUROK1



Neurokinin B (TAC3) (NM_013251) Human Recombinant Protein - TP720967XL

Summary:

This gene encodes a member of the tachykinin family of secreted neuropeptides. The encoded preproprotein is proteolytically processed to generate the mature peptide, which is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. This peptide is the ligand for the neurokinin-3 receptor. This protein is also expressed in the outer syncytiotrophoblast of the placenta and may be associated with pregnancy-induced hypertension and pre-eclampsia. Mutations in this gene are associated with normosmic hypogonadotropic hypogonadism. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016]

Protein Families:

Druggable Genome, Secreted Protein