

## Product datasheet for TP720974XL

## STAT6 (NM\_003153) Human Recombinant Protein

## **Product data:**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human signal transducer and activator of transcription 6, interleukin-4 induced (STAT6), transcript variant 2
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Ser627-Ser837
Tag:	C-His
Predicted MW:	23.9 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 $\mu m$ filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<u>NP 003144</u>
Locus ID:	6778
UniProt ID:	<u>P42226</u>
RefSeq Size:	4031
Cytogenetics:	12q13.3
RefSeq ORF:	2541
Synonyms:	D12S1644; IL-4-STAT; STAT6B; STAT6C



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	STAT6 (NM_003153) Human Recombinant Protein – TP720974XL
Summary:	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010]
Protein Families	: Druggable Genome, ES Cell Differentiation/IPS, Stem cell relevant signaling - DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway, Transcription Factors
Protein Pathway	<b>/s:</b> Jak-STAT signaling pathway

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US