

Product datasheet for TP720985M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NEDD8 (NM_006156) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human neural precursor cell expressed, developmentally

down-regulated 8 (NEDD8)

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Met1-Gly76

Tag: N-His&SUMO

Predicted MW: 20.9 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: <u>NP 006147</u>

Locus ID: 4738

UniProt ID: Q15843

RefSeq Size: 625

Cytogenetics: 14q12

RefSeq ORF: 243

Synonyms: NEDD-8





NEDD8 (NM_006156) Human Recombinant Protein – TP720985M

Summary: Ubiquitin-like protein which plays an important role in cell cycle control and embryogenesis.

> Covalent attachment to its substrates requires prior activation by the E1 complex UBE1C-APPBP1 and linkage to the E2 enzyme UBE2M. Attachment of NEDD8 to cullins activates their

associated E3 ubiquitin ligase activity, and thus promotes polyubiquitination and

proteasomal degradation of cyclins and other regulatory proteins.[UniProtKB/Swiss-Prot

Function]

Protein Families: Druggable Genome