

### Product datasheet for TP721030

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# IGF1 (NM\_000618) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human insulin-like growth factor 1 (somatomedin C) (IGF1),

transcript variant 4

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Gly49-Ala118

Tag:Tag FreePredicted MW:7.6 kDaConcentration:lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Lyophilized from a 0.2 um filtered solution of 20mM HAc-NaAc, pH 4.5.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not

recommended to reconstitute to a concentration less than 100  $\mu$ g/ml. Dissolve the lyophilized protein in 50mM Acetic Acid. Please aliquot the reconstituted solution to minimize freeze-

thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Stability:** Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 000609

**Locus ID:** 3479

UniProt ID: P05019

RefSeq Size: 7260

Cytogenetics: 12q23.2





#### IGF1 (NM\_000618) Human Recombinant Protein - TP721030

RefSeq ORF: 459

Synonyms: IGF; IGF-I; IGFI; MGF

**Summary:** The protein encoded by this gene is similar to insulin in function and structure and is a

member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing

to generate mature protein. [provided by RefSeq, Sep 2015]

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

Protein Pathways: Dilated cardiomyopathy, Focal adhesion, Glioma, Hypertrophic cardiomyopathy (HCM), Long-

term depression, Melanoma, mTOR signaling pathway, Oocyte meiosis, p53 signaling pathway, Pathways in cancer, Progesterone-mediated oocyte maturation, Prostate cancer