

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP721043

IDE (NM_004969) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human insulin-degrading enzyme (IDE), transcript variant 1
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Met42-Leu1019
Tag:	C-His
Predicted MW:	41.27 kDa
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Supplied as a 0.2 um filtered solution of 20mM Tris-HCl, 150mM NaCl, 0.05%Brij35, 10% Glycerol, pH 7.5.
Endotoxin:	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
Storage:	Store at < -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.
Stability:	Stable for at least 3 months from date of receipt under proper storage and handling conditions.
RefSeq:	<u>NP 004960</u>
Locus ID:	3416
UniProt ID:	<u>P14735</u>
RefSeq Size:	3279
Cytogenetics:	10q23.33
RefSeq ORF:	3057
Synonyms:	INSULYSIN



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE	IDE (NM_004969) Human Recombinant Protein – TP721043	
---------	------------------------------------------------------	--

Summary: This gene encodes a zinc metallopeptidase that degrades intracellular insulin, and thereby terminates insulins activity, as well as participating in intercellular peptide signalling by degrading diverse peptides such as glucagon, amylin, bradykinin, and kallidin. The preferential affinity of this enzyme for insulin results in insulin-mediated inhibition of the degradation of other peptides such as beta-amyloid. Deficiencies in this protein's function are associated with Alzheimer's disease and type 2 diabetes mellitus but mutations in this gene have not been shown to be causitive for these diseases. This protein localizes primarily to the cytoplasm but in some cell types localizes to the extracellular space, cell membrane, peroxisome, and mitochondrion. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described but have not been experimentally verified.[provided by RefSeq, Sep 2009]
Protein Families: Druggable Genome, Protease

Protein Pathways: Alzheimer's disease

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US