

Product datasheet for TP721048L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Syndecan 1 (SDC1) (NM 002997) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human syndecan 1 (SDC1), transcript variant 2

Species: Human **HEK293 Expression Host:**

Expression cDNA Clone

Tag:

or AA Sequence:

Predicted MW: 20.39 kDa **Concentration:** lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)

Gln23-Glu251

C-6His

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Store at -80°C. Storage:

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

NP 002988 RefSeq:

Locus ID: 6382 **UniProt ID:** P18827 RefSeg Size: 3217 Cytogenetics: 2p24.1 RefSeq ORF: 930

Synonyms: CD138; SDC; SYND1; syndecan





Syndecan 1 (SDC1) (NM_002997) Human Recombinant Protein - TP721048L

Summary:

The protein encoded by this gene is a transmembrane (type I) heparan sulfate proteoglycan and is a member of the syndecan proteoglycan family. The syndecans mediate cell binding, cell signaling, and cytoskeletal organization and syndecan receptors are required for internalization of the HIV-1 tat protein. The syndecan-1 protein functions as an integral membrane protein and participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. Altered syndecan-1 expression has been detected in several different tumor types. While several transcript variants may exist for this gene, the full-length natures of only two have been described to date. These two represent the major variants of this gene and encode the same protein. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), ECM-receptor interaction