

Product datasheet for **TP721052M**

IL20 Receptor alpha (IL20RA) (NM_014432) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human interleukin 20 receptor, alpha (IL20RA)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Val30-Lys250
Tag:	C-Fc
Predicted MW:	26.3 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	NP_055247
Locus ID:	53832
UniProt ID:	Q9UHF4
RefSeq Size:	3701
Cytogenetics:	6q23.3
RefSeq ORF:	1659
Synonyms:	CRF2-8; IL-20R-alpha; IL-20R1; IL-20RA



[View online »](#)

Summary: This gene encodes a member of the type II cytokine receptor family. The encoded protein is a subunit of the receptor for interleukin 20, a cytokine that may be involved in epidermal function. The interleukin 20 receptor is a heterodimeric complex consisting of the encoded protein and interleukin 20 receptor beta. This gene and interleukin 20 receptor beta are highly expressed in skin, and are upregulated in psoriasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway