

## Product datasheet for TP721059XL

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Activin Receptor Type IIA (ACVR2A) (NM 001616) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human activin A receptor, type IIA (ACVR2A)

Species: Human Expression Host: HEK293

**Expression cDNA Clone** 

ne .

or AA Sequence:

Ala20-Pro134

Tag: C-Fc&His

Predicted MW: 41.2 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 001607

Locus ID: 92

 UniProt ID:
 P27037

 RefSeq Size:
 5244

Cytogenetics: 2q22.3-q23.1

RefSeq ORF: 1539

Synonyms: ACTRII; ACVR2





## Activin Receptor Type IIA (ACVR2A) (NM\_001616) Human Recombinant Protein - TP721059XL

**Summary:** 

This gene encodes a receptor that mediates the functions of activins, which are members of the transforming growth factor-beta (TGF-beta) superfamily involved in diverse biological processes. The encoded protein is a transmembrane serine-threonine kinase receptor which mediates signaling by forming heterodimeric complexes with various combinations of type I and type II receptors and ligands in a cell-specific manner. The encoded type II receptor is primarily involved in ligand-binding and includes an extracellular ligand-binding domain, a transmembrane domain and a cytoplasmic serine-threonine kinase domain. This gene may be associated with susceptibility to preeclampsia, a pregnancy-related disease which can result in maternal and fetal morbidity and mortality. Alternative splicing results in multiple

transcript variants of this gene. [provided by RefSeq, Jun 2013]

**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, TGF-beta signaling pathway