

## Product datasheet for **TP721068L**

### CD16b (FCGR3B) (NM\_000570) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human Fc fragment of IgG, low affinity IIIb, receptor (CD16b) (FCGR3B)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Thr20-Gln208
Tag:	C-Fc&His
Predicted MW:	49.3 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<a href="#">NP_000561</a>
Locus ID:	2215
UniProt ID:	<a href="#">O75015</a> , <a href="#">A0A3B3ISU3</a>
RefSeq Size:	2148
Cytogenetics:	1q23.3
RefSeq ORF:	699
Synonyms:	CD16; CD16A; CD16b; FCG3; FCGR3; FCGR3A; FCR-10; FCRIII; FCRIIIb



[View online »](#)

<b>Summary:</b>	The protein encoded by this gene is a low affinity receptor for the Fc region of gamma immunoglobulins (IgG). The encoded protein acts as a monomer and can bind either monomeric or aggregated IgG. This gene may function to capture immune complexes in the peripheral circulation. Several transcript variants encoding different isoforms have been found for this gene. A highly-similar gene encoding a related protein is also found on chromosome 1. [provided by RefSeq, Aug 2012]
<b>Protein Families:</b>	ES Cell Differentiation/IPS, Secreted Protein, Transmembrane
<b>Protein Pathways:</b>	Natural killer cell mediated cytotoxicity, Systemic lupus erythematosus