

# **Product datasheet for TP721072M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NrCAM (NM\_005010) Human Recombinant Protein

### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human neuronal cell adhesion molecule (NRCAM), transcript

variant 2

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Gln25-Asn600

Tag: C-Fc

Predicted MW: 91.2 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

**Stability:** Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** <u>NP 005001</u>

**Locus ID:** 4897

UniProt ID: Q92823

RefSeq Size: 6322

Cytogenetics: 7q31.1

RefSeq ORF: 3549





#### NrCAM (NM\_005010) Human Recombinant Protein - TP721072M

**Summary:** 

Cell adhesion molecules (CAMs) are members of the immunoglobulin superfamily. This gene encodes a neuronal cell adhesion molecule with multiple immunoglobulin-like C2-type domains and fibronectin type-III domains. This ankyrin-binding protein is involved in neuron-neuron adhesion and promotes directional signaling during axonal cone growth. This gene is also expressed in non-neural tissues and may play a general role in cell-cell communication via signaling from its intracellular domain to the actin cytoskeleton during directional cell migration. Allelic variants of this gene have been associated with autism and addiction vulnerability. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs)