

Product datasheet for TP721083L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eotaxin 2 (CCL24) (NM 002991) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human chemokine (C-C motif) ligand 24 (CCL24)

Species: Human Expression Host: HEK293

Expression cDNA Clone

Val27-Cys119

or AA Sequence:

Tag: C-His

Predicted MW: 11.5 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 002982

 Locus ID:
 6369

 UniProt ID:
 000175

 RefSeq Size:
 360

 Cytogenetics:
 7q11.23

RefSeq ORF: 357

Synonyms: Ckb-6; MPIF-2; MPIF2; SCYA24





Eotaxin 2 (CCL24) (NM_002991) Human Recombinant Protein - TP721083L

Summary:

This gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity on resting T lymphocytes, a minimal activity on neutrophils, and is negative on monocytes and activated T lymphocytes. This protein also has antimicrobial activity, displaying an antibacterial effect on S. pneumoniae, S. aureus, Nontypeable H. influenzae, and P. aeruginosa. Finally, the protein is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line. [provided by RefSeq, Jul 2020]

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction