

# Product datasheet for TP721100M

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LDL Receptor (LDLR) (NM\_000527) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human low density lipoprotein receptor (LDLR), transcript

variant 1

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Ala22-Arg788

Tag: C-AVI&His Predicted MW: 88.4 kDa

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeg: NP 000518

**Locus ID:** 3949

**UniProt ID:** <u>P01130</u>, <u>A0A024R7D5</u>

**RefSeq Size:** 5175

Cytogenetics: 19p13.2

RefSeq ORF: 2580

**Synonyms:** FH; FHC; FHCL1; LDLCQ2





### LDL Receptor (LDLR) (NM\_000527) Human Recombinant Protein - TP721100M

Summary: The low density lipoprotein receptor (LDLR) gene family consists of cell surface proteins

involved in receptor-mediated endocytosis of specific ligands. Low density lipoprotein (LDL) is normally bound at the cell membrane and taken into the cell ending up in lysosomes where the protein is degraded and the cholesterol is made available for repression of microsomal enzyme 3-hydroxy-3-methylglutaryl coenzyme A (HMG CoA) reductase, the rate-limiting step in cholesterol synthesis. At the same time, a reciprocal stimulation of cholesterol ester synthesis takes place. Mutations in this gene cause the autosomal dominant disorder, familial hypercholesterolemia. Alternate splicing results in multiple transcript variants.[provided by

RefSeq, Sep 2010]

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

**Protein Pathways:** Endocytosis