

Product datasheet for TP721102M

CFHR5 (NM_030787) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Purified recombinant protein of Human complement factor H-related 5 (CFHR5) Species: Human **Expression Host: HEK293** Glu19-Glu569 Expression cDNA Clone or AA Sequence: C-His Tag: Predicted MW: 63.5 kDa **Concentration:** lot specific **Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Endotoxin: Endotoxin level is < 0.1 ng/ μ g of protein (< 1 EU/ μ g) Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 Storage: weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Stable for at least 6 months from date of receipt under proper storage and handling Stability: conditions. NP 110414 **RefSeq:** 81494 Locus ID: **UniProt ID:** Q9BXR6 2823 **RefSeq Size:** Cytogenetics: 1q31.3 **RefSeq ORF:** 1707 Synonyms: CFHL5; CFHR5D; FHR-5; FHR5



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	CFHR5 (NM_030787) Human Recombinant Protein – TP721102M
Summary:	This gene is a member of a small complement factor H (CFH) gene cluster on chromosome 1. Each member of this gene family contains multiple short consensus repeats (SCRs) typical of regulators of complement activation. The protein encoded by this gene has nine SCRs with the first two repeats having heparin binding properties, a region within repeats 5-7 having heparin binding and C reactive protein binding properties, and the C-terminal repeats being similar to a complement component 3 b (C3b) binding domain. This protein co-localizes with C3, binds C3b in a dose-dependent manner, and is recruited to tissues damaged by C- reactive protein. Allelic variations in this gene have been associated, but not causally linked, with two different forms of kidney disease: membranoproliferative glomerulonephritis type II (MPGNII) and hemolytic uraemic syndrome (HUS). [provided by RefSeq, Jan 2010]
Protein Families	s: Secreted Protein, Transmembrane

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US