

Product datasheet for TP721109XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Chromogranin C (SCG2) (NM_003469) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human secretogranin II (SCG2)

Species: Human Expression Host: HEK293

Expression cDNA Clone

Ser29-Met617

or AA Sequence:

Tag: C-His

Predicted MW: 68.8 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 003460

 Locus ID:
 7857

 UniProt ID:
 P13521

 RefSeq Size:
 2586

 Cytogenetics:
 2q36.1

 RefSeq ORF:
 1851

Synonyms: CHGC; EM66; SgII; SN





Chromogranin C (SCG2) (NM_003469) Human Recombinant Protein - TP721109XL

Summary: The protein encoded by this gene is a member of the chromogranin/secretogranin family of

neuroendocrine secretory proteins. Studies in rodents suggest that the full-length protein,

secretogranin II, is involved in the packaging or sorting of peptide hormones and

neuropeptides into secretory vesicles. The full-length protein is cleaved to produce the active peptide secretoneurin, which exerts chemotaxic effects on specific cell types, and EM66,

whose function is unknown. [provided by RefSeq, Jul 2008]

Protein Families: Secreted Protein, Transmembrane