

## **Product datasheet for TP721158XL**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Adipoq (NM\_009605) Mouse Recombinant Protein

## **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse adiponectin, C1Q and collagen domain containing

(Adipoq)

Species: Mouse Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Glu18-Asn247

Tag: N-His

Predicted MW: 27.2 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 033735

 Locus ID:
 11450

 UniProt ID:
 Q60994

 RefSeq Size:
 1233

Cytogenetics: 16 13.96 cM

RefSeg ORF: 741

Synonyms: 30kDa; Acdc; Acrp30; Ad; adipo; apM1; APN; GBP28







**Summary:** 

Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW.[UniProtKB/Swiss-Prot Function]