

# Product datasheet for TP721175M

## IL1 beta (IL1B) (NM\_000576) Human Recombinant Protein

### **Product data:**

#### **Product Type: Recombinant Proteins Description:** Purified recombinant protein of Human interleukin 1, beta (IL1B) Species: Human **Expression Host:** E. coli **Expression cDNA Clone** Met1-Leu586 or AA Sequence: Tag Free Tag: **Predicted MW:** 17.4 kDa **Concentration:** lot specific **Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl Endotoxin: Endotoxin level is < 0.1 ng/ $\mu$ g of protein (< 1 EU/ $\mu$ g) Store at -80°C. Storage: Stable for at least 6 months from date of receipt under proper storage and handling Stability: conditions. **RefSeq:** NP 000567 Locus ID: 3553 **UniProt ID:** P01584 **RefSeq Size:** 1498 Cytogenetics: 2q14.1 **RefSeq ORF:** 807 Synonyms: IL-1; IL1-BETA; IL1beta; IL1F2



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	IL1 beta (IL1B) (NM_000576) Human Recombinant Protein – TP721175M
Summary:	The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly, IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of pro-inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2020]
Protein Families	: Druggable Genome, Secreted Protein
Protein Pathway	<ul> <li>Alzheimer's disease, Apoptosis, Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, Graft-versus-host disease, Hematopoietic cell lineage, MAPK signaling pathway, NOD-like receptor signaling pathway, Prion diseases, Toll-like receptor signaling pathway, Type I diabetes mellitus</li> </ul>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US