

## Product datasheet for **TP721198M**

### HMG1 (HMGB1) (NM\_002128) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human high mobility group box 1 (HMGB1)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Pro92-Val176
Tag:	N-MARI
Predicted MW:	10 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<a href="#">NP_002119</a>
Locus ID:	3146
UniProt ID:	<a href="#">P09429</a> , <a href="#">A0A024RDR0</a> , <a href="#">Q5T7C3</a>
RefSeq Size:	3428
Cytogenetics:	13q12.3
RefSeq ORF:	645
Synonyms:	HMG-1; HMG1; HMG3; SBP-1



[View online »](#)

**Summary:**

This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2015]

**Protein Families:**

Druggable Genome, Stem cell - Pluripotency, Transcription Factors

**Protein Pathways:**

Base excision repair