

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP721210XL

PDGF AA (PDGFA) (NM_033023) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human platelet-derived growth factor alpha polypeptide (PDGFA), transcript variant 2
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Ser87-Thr211
Tag:	Tag Free
Predicted MW:	14.1 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Reconstitution Method: Storage:	recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized
	recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling
Storage: Stability:	recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions.
Storage: Stability: RefSeq:	recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 148983</u>
Storage: Stability: RefSeq: Locus ID:	recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. NP 148983 5154
Storage: Stability: RefSeq: Locus ID: UniProt ID:	recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 148983</u> 5154 <u>P04085</u>
Storage: Stability: RefSeq: Locus ID: UniProt ID: RefSeq Size:	recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 4mM HCl . Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 148983</u> 5154 <u>P04085</u> 2749



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PDGF AA (PDGFA) (NM_033023) Human Recombinant Protein – TP721210XL
Summary:	This gene encodes a member of the protein family comprised of both platelet-derived growth factors (PDGF) and vascular endothelial growth factors (VEGF). The encoded preproprotein is proteolytically processed to generate platelet-derived growth factor subunit A, which can homodimerize, or alternatively, heterodimerize with the related platelet-derived growth factor subunit B. These proteins bind and activate PDGF receptor tyrosine kinases, which play a role in a wide range of developmental processes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2015]
Protein Families:	Druggable Genome
Protein Pathways	: Cytokine-cytokine receptor interaction, Focal adhesion, Gap junction, Glioma, MAPK signaling pathway, Melanoma, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US