

Product datasheet for TP721232L

LRRC25 (NM_145256) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Purified recombinant protein of Human leucine rich repeat containing 25 (LRRC25) Species: Human **HEK293 Expression Host:** Leu21-Thr165 **Expression cDNA Clone** or AA Sequence: C-His Tag: Predicted MW: 16.7 kDa **Concentration:** lot specific **Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Endotoxin: Endotoxin level is < 0.1 ng/ μ g of protein (< 1 EU/ μ g) **Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Lyophilized protein should be stored at $< -20^{\circ}$ C, though stable at room temperature for 3 Storage: weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Stability: Stable for at least 6 months from date of receipt under proper storage and handling conditions. **RefSeq:** NP 660299 Locus ID: 126364 UniProt ID: Q8N386 **RefSeq Size:** 2430 **Cytogenetics:** 19p13.11 **RefSeq ORF:** 915 MAPA Synonyms:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	LRRC25 (NM_145256) Human Recombinant Protein – TP721232L
Summary:	Plays a role in the inhibition of RLR-mediated type I interferon signaling pathway by targeting DDX58/RIG-I for autophagic degradation. Interacts specifically with ISG15-associated DDX58 to promote interaction between DDX58 and the autophagic cargo receptor p62/SQSTM1 to mediate DDX58 degradation via selective autophagy (PubMed:29288164). Plays also a role in the inhibition of NF-kappa-B signaling pathway and inflammatory response by promoting the degradation of p65/RELA.[UniProtKB/Swiss-Prot Function]
Protein Familie	rs: Transmembrane

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US