

# **Product datasheet for TP721232XL**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LRRC25 (NM\_145256) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human leucine rich repeat containing 25 (LRRC25)

Species: Human
Expression Host: HEK293

**Expression cDNA Clone** 

Leu21-Thr165

or AA Sequence:

Tag: C-His

Predicted MW: 16.7 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 660299

 Locus ID:
 126364

 UniProt ID:
 Q8N386

 RefSeq Size:
 2430

Cytogenetics: 19p13.11

**RefSeq ORF:** 915 **Synonyms:** MAPA





### LRRC25 (NM\_145256) Human Recombinant Protein - TP721232XL

Summary: Plays a role in the inhibition of RLR-mediated type I interferon signaling pathway by targeting

DDX58/RIG-I for autophagic degradation. Interacts specifically with ISG15-associated DDX58 to promote interaction between DDX58 and the autophagic cargo receptor p62/SQSTM1 to mediate DDX58 degradation via selective autophagy (PubMed:29288164). Plays also a role in the inhibition of NF-kappa-B signaling pathway and inflammatory response by promoting the

degradation of p65/RELA.[UniProtKB/Swiss-Prot Function]

**Protein Families:** Transmembrane