

## Product datasheet for TP721241L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL5 (NM\_000879) Human Recombinant Protein

## **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human interleukin 5 (colony-stimulating factor, eosinophil)

(IL5)

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

lle20-Ser134

Tag: C-His

Predicted MW: 14.2 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

**Stability:** Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** <u>NP 000870</u>

 Locus ID:
 3567

 UniProt ID:
 P05113

 RefSeq Size:
 816

Cytogenetics: 5q31.1 RefSeq ORF: 402

Synonyms: EDF; IL-5; TRF





**Summary:** 

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2. This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq, Jul 2013]

**Protein Families:** 

Druggable Genome, Secreted Protein

**Protein Pathways:** 

Allograft rejection, Asthma, Autoimmune thyroid disease, Cytokine-cytokine receptor interaction, Fc epsilon RI signaling pathway, Hematopoietic cell lineage, Jak-STAT signaling pathway, T cell receptor signaling pathway