

Product datasheet for TP721253

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FCGR1A Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: APC Conjugated Human CD64 Protein (C-His)

Species: Human Expression Host: CHO

Expression cDNA Clone

Gln16-Thr287

or AA Sequence:

Tag: C-His

Predicted MW: The protein has the predicted molecular weight of 32kDa and migrates at approximately 50-

60 kDa on SDS-PAGE with DTT-reduced condition before APC conjugatoin.

Concentration: 25µg size is bottled at 0.1mg/mL concentration. 100 µg size is bottled at lot specific

concentration.

Purity: >90%
Conjugation: APC

Buffer: 1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein

Bioactivity: Positive

The definition of the active protein (purified and biotinylated) is defined as the protein that can bind to its biological receptor/ligand. For conjugated protein, it is defined with its function

to bind to the ScFv of the active CAR-transfected cells in flow cytometry test.

Preparation: Affinity Ni-NTA

Applications: FACS

Storage: An unopened vial can be stored at 4°C for 2 weeks or at -20°C and below for six months. This

stock solution should be aliquoted and stored at \leq -70°C to minimize the freeze/thaw cycles.

Stability: 6 Months

RefSeq: <u>NP 000557.1</u>

Locus ID: 2209 UniProt ID: P12314





FCGR1A Human Recombinant Protein - TP721253

Summary: Receptors for the Fc portion of human IgG (Fc gamma receptors) can be divided into three

classes as Fc gamma RI (CD64), RII (CD32), and RIII (CD16). Each of these proteins may be coded by multiple genes and exist in different isoforms. Fc gamma RI or CD64 proteins can bind Fc portion of human IgG with high affinity and play important role in early immune response. In contrast CD16 and CD32 bind human IgG with low affinity and can only recognize the IgG aggregates with multivalent antigens. Human CD64 is expressed

constitutively on macrophages and monocytes and its expression can be induced by IFNg or

G-CSF on polymorphonuclear leukocytes.

Protein Families: FACS

Protein Pathways: Fc gamma R-mediated phagocytosis, Hematopoietic cell lineage, Systemic lupus

erythematosus