

## Product datasheet for TP721362

## **SECTM1 Human Recombinant Protein**

**Product data:** 

**Product Type: Recombinant Proteins** 

Description: PE Conjugated Human SECTM1 Protein (C-His)

Species: Human CHO **Expression Host:** 

**Expression cDNA Clone** 

Gln29-Gly145

or AA Sequence:

Tag:

C-His

**Predicted MW:** The protein has a predicted molecular weight of 14kDa and migrates at approximately 20kDa

on SDS-PAGE with DTT-reduced condition before PE conjugation.

Concentration: 25μg size is bottled at 0.1mg/mL concentration. 100 μg size is bottled at lot specific

concentration.

>90% **Purity:** 

Conjugation: PF

**Buffer:** 1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein

**Bioactivity:** Positive

> The definition of the active protein (purified and biotinylated) is defined as the protein that can bind to its biological receptor/ligand. For conjugated protein, it is defined with its function

to bind to the ScFv of the active CAR-transfected cells in flow cytometry test.

Preparation: Affinity Ni-NTA

**Applications: FACS** 

Storage: An unopened vial can be stored at 4°C for 2 weeks or at -20°C and below for six months. This

stock solution should be aliquoted and stored at  $\leq$  -70°C to minimize the freeze/thaw cycles.

Stability: 6 Months RefSeq: Q8WVN6

Locus ID: 6398

UniProt ID: Q8WVN6



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Summary:

Human SECTM1 (secreted and transmembrance 1) is a type I transmembrane glycoprotein consisted of a 28 aa signal sequence, a 117 aa extracellular domain (ECD) with one potential N-linked glycosylation site, a 21 aa transmembrane sequence, and an 82 aa cytoplasmic sequence. It is localized in the Golgi apparatus and exists as a transmembrane and a soluble form. SECTM1 protein expression is detected in some myeloid cells, such as stimulated monocytes, immature monocyte-derived dendritic cells, and granulocytes. In the thymus, it is also expressed by epithelia and fibroblasts. Its expression is stimulated by IFN-gamma. Although its function is not well understood, the soluble N-terminus form of human SECTM1 demonstrates that it is a costimulatory ligand for human CD4 and CD8 T cell proliferation and cytokine production in CD7-dependent manner.

Protein Families:

**FACS** 

## **Product images:**



PE conjugated Human SECTM1 Protein (C-His) on SDS-PAGE under reducing condition PE (+) and non-reducing condition PE(-). C(-) is unconjugated Human SECTM1 Protein under non reducing condition.