

## **Product datasheet for TP723120**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Adipoq (NM\_009605) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse adiponectin, C1Q and collagen domain containing

(Adipoq).

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MKGEPGEAAY MYRSAFSVGL ETRVTVPNVP IRFTKIFYNQ QNHYDGSTGK FYCNIPGLYY FSYHITVYMK DVKVSLFKKD KAVLFTYDQY QEKNVDQASG SVLLHLEVGD QVWLQVYGDG

DHNGLYADNV NDSTFTGFLL YHDTN

Tag: Tag Free
Predicted MW: 16.6 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 μM filtered solution of 20mM phosphate buffer,100mM NaCl, pH 7.2

**Bioactivity:** Determined by its ability to inhibit the proliferation of murine myeloid cell lines M1. ED50 for

this effect is less than or equal to 15 ug/ml.

**Endotoxin:** Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)

Storage: Store at -80°C.

**Stability:** Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 033735

 Locus ID:
 11450

 UniProt ID:
 Q60994

 RefSeq Size:
 1233

Cytogenetics: 16 13.96 cM

RefSeg ORF: 741

Synonyms: 30kDa; Acdc; Acrp30; Ad; adipo; apM1; APN; GBP28







**Summary:** 

Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW.[UniProtKB/Swiss-Prot Function]