

## **Product datasheet for TP723274**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cxcl5 (NM\_009141) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse chemokine (C-X-C motif) ligand 5 (Cxcl5).

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

TELRCVCLTV TPKINPKLIA NLEVIPAGPQ CPTVEVIAKL KNQKEVCLDP EAPVIKKIIQ KILGSDKKKA

Tag: Tag Free
Predicted MW: 7.6 kDa
Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 μM filtered solution of 20mM phosphate buffer,100mM NaCl, pH 7.2

**Bioactivity:** Determined by its ability to chemoattract human neutrophils using a concentration range of

10.0-100.0 ng/ml.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 033167

 Locus ID:
 20311

 UniProt ID:
 P50228

 RefSeq Size:
 1655

Cytogenetics: 5 44.78 cM

RefSeg ORF: 396

Synonyms: AMCF; AMCF-II; Cxcl; Cxcl6; ENA-; ENA-78; GCP-2; L; LIX; Scyb; Scyb5; Scyb6





**Summary:** 

This gene encodes a protein that is a member of the CXC subfamily of chemokines. Chemokines, which recruit and activate leukocytes, are classified by function (inflammatory or homeostatic) or by structure. This protein is proposed to bind the G-protein coupled receptor chemokine (C-X-C motif) receptor 2 to recruit neutrophils and to have homeostatic and inflammatory functions. In mouse, deficiency of this gene is associated with increased lung inflammation that is neutrophil-dependent. [provided by RefSeq, May 2013]

## **Product images:**

