

Product datasheet for TP723282

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MCP2 (CCL8) (NM_005623) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human chemokine (C-C motif) ligand 8 (CCL8).

Species: Human
Expression Host: E. coli

Expression cDNA Clone QPDSVSIPIT CCFNVINRKI PIQRLESYTR ITNIQCPKEA VIFKTKRGKE VCADPKERWV RDSMKHLDQI

or AA Sequence: FQNLKP

Tag: Tag Free
Predicted MW: 8.9 kDa
Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 μM filtered solution of 20mM phosphate buffer,100mM NaCl, pH 7.2

Bioactivity: Determined by its ability to chemoattract human peripheral blood monocytes using a

concentration range of 10.0-100.0 ng/ml.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 005614

 Locus ID:
 6355

 UniProt ID:
 P80075

 RefSeq Size:
 1351

 Cytogenetics:
 17q12

 RefSeq ORF:
 297

Synonyms: HC14; MCP-2; MCP2; SCYA8; SCYA10





Summary:

This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils. By recruiting leukocytes to sites of inflammation this cytokine may contribute to tumor-associated leukocyte infiltration and to the antiviral state against HIV infection. [provided by RefSeq, Sep 2014]

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction, NOD-like receptor

signaling pathway

Product images:

