

## Product datasheet for **TP723354**

### **Pdgfa (NM\_008808) Mouse Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse platelet derived growth factor, PDGFAB, a disulfide-linked dimer, consisting of one A chain and one B chain.
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	<b>alphachain</b> MSIEEAVPAV CKTRTVIYEI PRSQVDPTSA NFLIWPPCVE VKRCTGCCNT SSVKQCPSRV HHRSVKVAKV EYVRKKPKLK EVQVRLEEHL ECACATTSLN PDYREEDTGR PRESGKKRKR KRLKPT <b>betachain</b> MSLGSLTIAE PAMIAECKTR TEVFEISRRL IDRTNANFLV WPPCVEVQRC SGCCNNRNVQ CRPTQVQLRP VQVRKIGIVR KKPIFKKATV TLGDHLACKC ETVAAARPVT
Tag:	Tag Free
Predicted MW:	25.5 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Lyophilized from a 0.2 $\mu$ M filtered solution of 20mM phosphate buffer, 100mM NaCl, pH 7.2
Bioactivity:	ED50 as determined by the dose-dependent stimulation of thymidine uptake by BALB/c 3T3 cells is less than or equal to 1 ng/ml, corresponding to a specific activity of $> 1 \times 10^6$ units/mg.
Endotoxin:	Endotoxin level is $< 0.1$ ng/ $\mu$ g of protein ( $< 1$ EU/ $\mu$ g)
Storage:	Store at $-80^{\circ}\text{C}$ .
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<a href="#">NP_032834</a>
Locus ID:	18590
UniProt ID:	<a href="#">P20033</a>
RefSeq Size:	1019
Cytogenetics:	5 77.65 cM
RefSeq ORF:	588



[View online »](#)

**Summary:**

Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells of mesenchymal origin. Required for normal lung alveolar septum formation during embryogenesis, normal development of the gastrointestinal tract, normal development of Leydig cells and spermatogenesis. Required for normal oligodendrocyte development and normal myelination in the spinal cord and cerebellum. Plays an important role in wound healing. Signaling is modulated by the formation of heterodimers with PDGFB. [UniProtKB/Swiss-Prot Function]

**Product images:**