

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TP723460

## DR5 (TNFRSF10B) (NM\_003842) Human Recombinant Protein

## **Product data:**

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human tumor necrosis factor receptor superfamily, member 10b (TNFRSF10B), transcript variant 1.
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MESALITQQD LAPQQRVAPQ QKRSSPSEGL CPPGHHISED GRDCISCKYG QDYSTHWNDL LFCLRCTRCD SGEVELSPCT TTRNTVCQCE EGTFREEDSP EMCRKCRTGC PRGMVKVGDC TPWSDIECVH KES
Tag:	Tag Free
Predicted MW:	14.9 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Lyophilized from a 0.2 $\mu$ M filtered solution of 20mM phosphate buffer,100mM NaCl, pH 7.2
Bioactivity:	sTRAIL Rec 2 reduced the production of LPS-induced TNF by its ability to neutralize endogenous TRAIL in fresh human PBMC. In this assay, endogenous TRAIL is induced during a 24 hour exposure to LPS (10 ng/mL) but in the presence ofsTRAIL Rec 2, TRAIL-induced TNF is suppressed.
Endotoxin:	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<u>NP 671716</u>
Locus ID:	8795
UniProt ID:	<u>O14763</u> , <u>Q7Z2I8</u>
RefSeq Size:	4154
Cytogenetics:	8p21.3
RefSeq ORF:	1320



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>DR5 (TNFRSF10B) (NM_003842) Human Recombinant Protein – TP723460</b>	
Synonyms:	CD262; DR5; KILLER; KILLER/DR5; TRAIL-R2; TRAILR2; TRICK2; TRICK2A; TRICK2B; TRICKB; ZTNFR9
Summary:	The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene. [provided by RefSeq, Mar 2009]
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Apoptosis, Cytokine-cytokine receptor interaction, Natural killer cell mediated cytotoxicity, p53 signaling pathway

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US