

Product datasheet for **TP723727**

Cxcl10 (NM_021274) Mouse Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse chemokine (C-X-C motif) ligand 10 (Cxcl10 / IP10)
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Mouse CXCL10, the region of Ile22-Pro98, from gene Accession# NM_021274
Tag:	Tag Free
Predicted MW:	8.8 kDa
Concentration:	lot specific
Purity:	>98%, as determined by Coomassie stained SDS-PAGE.
Buffer:	1 x PBS
Bioactivity:	Recombinant Mouse CXCL10 (IP-10) induces chemotaxis of mouse Baf3-mCXCR3 transfectant cells in a dose-dependent manner. The ED50 for this effect is < 6 ng /mL.
Endotoxin:	Less than 0.01 ng per µg protein as determined by the LAL method
Storage:	Store at -80°C.
Stability:	Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6 months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated freeze/thaw cycles.
RefSeq:	NP_067249
Locus ID:	15945
UniProt ID:	P17515 , Q548V9
RefSeq Size:	1120
Cytogenetics:	5 46.57 cM
RefSeq ORF:	297
Synonyms:	C7; CRG-2; gIP-10; Ifi10; INP10; IP-10; IP10; mob-1; Scyb10



[View online »](#)

Summary:

Pro-inflammatory cytokine that is involved in a wide variety of processes such as chemotaxis, differentiation, and activation of peripheral immune cells, regulation of cell growth, apoptosis and modulation of angiostatic effects (By similarity) (PubMed:28623423). Plays thereby an important role during viral infections by stimulating the activation and migration of immune cells to the infected sites (PubMed:18624292, PubMed:19017990, PubMed:28468883). Mechanistically, binding of CXCL10 to the CXCR3 receptor activates G protein-mediated signaling and results in downstream activation of phospholipase C-dependent pathway, an increase in intracellular calcium production and actin reorganization. In turn, recruitment of activated Th1 lymphocytes occurs at sites of inflammation (By similarity). Activation of the CXCL10/CXCR3 axis plays also an important role in neurons in response to brain injury for activating microglia, the resident macrophage population of the central nervous system, and directing them to the lesion site. This recruitment is an essential element for neuronal reorganization (PubMed:15456824).[UniProtKB/Swiss-Prot Function]

Product images:

Recombinant Mouse CXCL10 (IP-10) induces chemotaxis of mouse Baf3-mCXCR3 transfectant cells in a dose-dependent manner. The ED50 for this effect is < 6 ng/mL.