

## **Product datasheet for TP723768**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Ccl2 (NM\_011333) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse chemokine (C-C motif) ligand 2 (Ccl2 / MCP-1)

Species: Mouse Expression Host: HEK293

**Expression cDNA Clone** 

or AA Sequence:

Tag: Tag Free

Predicted MW: 8.5 kDa

Concentration: lot specific

**Purity:** >98%, as determined by Coomassie stained SDS-PAGE.

**Buffer:** 1 x PBS

**Bioactivity:** The ED50 is 8 -15 ng/ml, corresponding to a specific activity of 0.6 - 1.25 x 10^5 units/mg,

determined by the dose dependent chemoattraction of THP-1 cells.

Mouse MCP1, the region of Gln24-Arg96, from gene Accession# NM\_011333

**Endotoxin:** Less than 0.01 ng per µg protein as determined by the LAL method

Storage: Store at -80°C.

Stability: Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6

months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated

freeze/thaw cycles.

**RefSeq:** <u>NP 035463</u>

 Locus ID:
 20296

 UniProt ID:
 P10148

 RefSeq Size:
 806

Cytogenetics: 11 49.82 cM

RefSeg ORF: 444

Synonyms: Al323594; HC11; JE; MCA; MCAF; MCP; MCP-; MCP-1; MCP1; Scy; Scya2; Sig; Sigje; SMC-C; SMC-

CF





**Summary:** 

This gene is one of several cytokine genes clustered on chromosome 11. Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and memory T cells but not for neutrophils. The human ortholog has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, such as psoriasis, rheumatoid arthritis, and atherosclerosis. [provided by RefSeq, Sep 2015]

## **Product images:**

