

Product datasheet for **TP723783**

Vegfa (NM_009505) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse vascular endothelial growth factor A (Vegfa), transcript variant 2
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Mouse VEGF120, the region of Ala27-Arg146, from gene Accession# NM_009505
Tag:	Tag Free
Predicted MW:	14.1 kDa
Concentration:	lot specific
Purity:	>98%, as determined by Coomassie stained SDS-PAGE.
Buffer:	5 mM Citric Acid, 5 mM NaH ₂ PO ₄ , 150 mM NaCl, pH 4.0
Bioactivity:	The ED ₅₀ is 1-4 ng/ml, corresponding to a specific activity of 1-0.25 x 10 ⁶ units/mg, determined by the dose dependent stimulation of HUVEC cells proliferation.
Endotoxin:	Less than 0.1 EU/μg (<0.01 ng/μg) protein as determined by the LAL method
Storage:	Store at -80°C.
Stability:	Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6 months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated freeze/thaw cycles.
RefSeq:	NP_033531
Locus ID:	22339
UniProt ID:	Q00731
RefSeq Size:	3475
Cytogenetics:	17 22.79 cM
RefSeq ORF:	1107
Synonyms:	V; Veg; Vegf; VEGF12; VEGF16; VEGF18; Vpf



[View online »](#)

Summary:

This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site.[provided by RefSeq, Nov 2015]

Product images: