

Product datasheet for **TP723788**

MCP2 (CCL8) (NM_005623) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human chemokine (C-C motif) ligand 8 (CCL8 / MCP-2)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Human CCL8, the region of Gln24-Pro99, from gene Accession# NM_005623.2
Tag:	Tag Free
Predicted MW:	8.9 kDa
Concentration:	lot specific
Purity:	>98%, as determined by Coomassie stained SDS-PAGE.
Buffer:	1 x PBS
Bioactivity:	Bioactivity was measured by its property to chemoattract human THP-1 cells in a dose dependent manner.
Endotoxin:	Less than 0.01 ng per µg protein as determined by the LAL method
Storage:	Store at -80°C.
Stability:	Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6 months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated freeze/thaw cycles.
RefSeq:	NP_005614
Locus ID:	6355
UniProt ID:	P80075
RefSeq Size:	1351
Cytogenetics:	17q12
RefSeq ORF:	297
Synonyms:	HC14; MCP-2; MCP2; SCYA8; SCYA10



[View online »](#)

Summary:

This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils. By recruiting leukocytes to sites of inflammation this cytokine may contribute to tumor-associated leukocyte infiltration and to the antiviral state against HIV infection. [provided by RefSeq, Sep 2014]

Protein Families:

Druggable Genome, Secreted Protein

Protein Pathways:

Chemokine signaling pathway, Cytokine-cytokine receptor interaction, NOD-like receptor signaling pathway