

Product datasheet for TP723794

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Exodus 2 (CCL21) (NM_002989) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human chemokine (C-C motif) ligand 21 (CCL21 / 6CKine)

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Human CCL21, the region of Ser24-Pro134, from gene Accession# NM_002989.2

Tag: Tag Free
Predicted MW: 12 kDa
Concentration: lot specific

Purity: >98%, as determined by Coomassie stained SDS-PAGE.

Buffer: 1 x PBS

Bioactivity: Bioactivity was measured by its property to chemoattract Baf3-hCCR7 transfectants in a dose

dependent manner.

Endotoxin: Less than 0.01 ng per µg protein as determined by the LAL method

Storage: Store at -80°C.

Stability: Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6

months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated

freeze/thaw cycles.

RefSeq: NP 002980

 Locus ID:
 6366

 UniProt ID:
 000585

 RefSeq Size:
 913

 Cytogenetics:
 9p13.3

RefSeq ORF: 402

Synonyms: 6Ckine; CKb9; ECL; SCYA21; SLC; TCA4





Exodus 2 (CCL21) (NM_002989) Human Recombinant Protein - TP723794

Summary:

This antimicrobial gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. Similar to other chemokines the protein encoded by this gene inhibits hemopoiesis and stimulates chemotaxis. This protein is chemotactic in vitro for thymocytes and activated T cells, but not for B cells, macrophages, or neutrophils. The cytokine encoded by this gene may also play a role in mediating homing of lymphocytes to secondary lymphoid organs. It is a high affinity functional ligand for chemokine receptor 7 that is expressed on T and B lymphocytes and a known receptor for another member of the cytokine family (small inducible cytokine A19). [provided by RefSeq, Sep 2014]

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction