

Product datasheet for TP723818

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eotaxin 3 (CCL26) (NM_006072) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human chemokine (C-C motif) ligand 26 (CCL26 Eotaxin-3)

Species: Human
Expression Host: E. coli

Expression cDNA Clone Human CCL26, the region of Thr24-Leu94, from gene Accession# NM_006072.4 was

or AA Sequence: expressed in E.coli.

Tag: Tag Free
Predicted MW: 8.4 kDa
Concentration: lot specific

Purity: >98%, as determined by Coomassie stained SDS-PAGE.

Buffer: 1 x PBS

Bioactivity: Bioactivity was measured by its property to chemoattract BaF3-mCCR3 transfectants in a

dose dependent manner.

Endotoxin: Less than 0.01 ng per µg protein as determined by the LAL method

Storage: Store at -80°C.

Stability: Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6

months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated

freeze/thaw cycles.

RefSeq: NP 006063

 Locus ID:
 10344

 UniProt ID:
 Q9Y258

 RefSeq Size:
 562

Cytogenetics: 7q11.23

RefSeg ORF: 282

Synonyms: IMAC; MIP-4a; MIP-4alpha; SCYA26; TSC-1





Summary:

This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. This protein also has antimicrobial activity, displaying an antibacterial effect on S. pneumoniae, S. aureus, Non-typeable H. influenzae, and P. aeruginosa. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil accumulation in atopic diseases. [provided by RefSeq, Jul 2020]

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction

Product images:

