

Product datasheet for TP723886

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TIMP2 (NM_003255) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human TIMP metallopeptidase inhibitor 2 (TIMP2)

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Human TIMP-2, the region of Cys27-Pro220, from gene Accession# NM_003255

Tag: C-His
Predicted MW: 22 kDa
Concentration: lot specific

Purity: >90%, as determined by Coomassie stained SDS-PAGE.

Buffer: 20 mM Tris, 150 mM NaCl, pH8.0

Bioactivity: Human TIMP-2 inhibits the activity of activated mouse MMP-9 (100 ng/mL). The is IC50 = 34

ng/mL.

Endotoxin: Less than 1.0 EU per μg of protein as determine by the LAL method

Storage: Store at -80°C.

Stability: Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6

months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated

freeze/thaw cycles.

RefSeq: NP 003246

Locus ID: 7077

UniProt ID: <u>P16035</u>, <u>A0A140VK57</u>

RefSeq Size: 3670 Cytogenetics: 17q25.3

RefSeq ORF: 660

Synonyms: CSC-21K; DDC8





Summary:

This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a result, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tissues undergoing remodelling of the extracellular matrix. [provided by RefSeq, Jul 2008]

Protein Families:

Druggable Genome, Secreted Protein

Product images:

