

## **Product datasheet for TP723980**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ACE2 Human Recombinant Protein (Angiotensin Converting Enzyme 2)**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Human ACE2 Protein, mFc Tag

Species: Human Expression Host: HEK293

**Expression cDNA Clone** 

or AA Sequence:

 $ACE2 \square Gln18 \hbox{-} Ser740 \square + mFc \square Pro99 \hbox{-} Lys330 \square$ 

Tag: C-Mouse Fc
Predicted MW: 109.7 kDa

**Purity:** > 95%

**Buffer:** Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants

before lyophilization

Reconstitution Method: Reconstitute with deionized water

**Preparation:** Affinity purification

**Applications:** ELISA

**Storage:** Store the lyophilized protein at -20°C.

After reconstitution, store the protein at -80°C for 12 months.

Avoid repeated freezing and thawing.

**Stability:** 12 months from date of despatch

 Locus ID:
 59272

 UniProt ID:
 Q9BYF1

**Summary:** The protein encoded by this gene belongs to the angiotensin-converting enzyme family of

dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-

specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-

NL63.





## **Product images:**




Figure 1. Human ACE2 Protein, mFc Tag on SDS-PAGE under reducing condition.

S-RBD, hFc Tagged protein ELISA 0.2 µg of S-RBD, hFc Tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) S-RBD, hFc tagged protein ([TP724020]) can bind Human ACE2, mFc Tagged proteinï¼TP723980ï¼ in a linear range of 0.976-39.35 ng/ml.