

# Product datasheet for TP724014

## **CTLA4 Human Recombinant Protein**

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins
21	
Description:	Human CTLA-4 Protein, hFc tag
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	CTLA-40Gln36-Ser1610+hFc0Glu99-Ala3300
Tag:	C-Human Fc
Predicted MW:	39.6 kDa
Purity:	> 95%
Buffer:	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization
<b>Reconstitution Method:</b>	Reconstitute with deionized water
Preparation:	Affinity purification
Applications:	Flow Cyt
Storage:	Store the lyophilized protein at -20°C. After reconstitution, store the protein at -80°C for 12 months. Avoid repeated freezing and thawing.
Stability:	12 months from date of despatch
Locus ID:	1493
UniProt ID:	<u>P16410</u>
Summary:	This gene is a member of the immunoglobulin superfamily and encodes a protein which transmits an inhibitory signal to T cells. The protein contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. Mutations in this gene have been associated with insulin-dependent diabetes mellitus, Graves disease, Hashimoto thyroiditis, celiac disease, systemic lupus erythematosus, thyroid-associated orbitopathy, and other autoimmune diseases.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



#### **Product images:**




Figure 1. Human CTLA-4 Protein, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human CTLA-4, hFc tagged protein (TP724014) can bind Anti-CTLA4 Neutralizing antibodyï¼BME100022ï¼ in a linear range of 0.64-80.0 ng/ml.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




Figure 3. HEK293 cell line transfected with irrelevant protein (B) and human B7-2 (A) were surface stained with Human CTLA4, hFc tagged protein (TP724014) 1 $\mu$ g/ml followed by Alexa 488-conjugated anti-human IgG secondary antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US