

## Product datasheet for **TP724065**

### Human TNFRSF11A Protein, hFc Tag

#### Product data:

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Human TNFRSF11A Protein, hFc Tag
<b>Expression Host:</b>	HEK293
<b>Tag:</b>	C-Human Fc
<b>Predicted MW:</b>	The protein has a predicted molecular mass of 46.2 kDa after removal of the signal peptide. The apparent molecular mass of TNFRSF11A-hFc is approximately 55-70 kDa due to glycosylation.
<b>Purity:</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Reconstitution Method:</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization.
<b>Storage:</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Stability:</b>	12 months from date of despatch
<b>Synonyms:</b>	Tumor necrosis factor receptor superfamily member 11A, ODFR, Receptor activator of NF-KB, CD265, RANK
<b>Summary:</b>	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptors can interact with various TRAF family proteins, through which this receptor induces the activation of NF-kappa B and MAPK8/JNK. This receptor and its ligand are important regulators of the interaction between T cells and dendritic cells. This receptor is also an essential mediator for osteoclast and lymph node development. Mutations at this locus have been associated with familial expansile osteolysis, autosomal recessive osteopetrosis, and Paget disease of bone. Alternatively spliced transcript variants have been described for this locus. [provided by RefSeq, Aug 2012]



[View online »](#)