

Product datasheet for TP724073

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Human GNRHR Protein, hFc Tag

Product data:

Product Type: Recombinant Proteins

Description: Human GNRHR Protein, hFc Tag

Expression Host: HEK293

Tag: C-Human Fc

Predicted MW: The protein has a predicted molecular mass of 30.2 kDa after removal of the signal

peptide. The apparent molecular mass of GNRHR-hFc is approximately 35-55 kDa due to

glycosylation.

Purity: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie

blue staining.

Reconstitution Method: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Stability: 12 months from date of despatch
Synonyms: GNRHR1; GRHR; HH7; LHRHR; LRHR

Summary: This gene encodes the receptor for type 1 gonadotropin-releasing hormone. This receptor is

a member of the seven-transmembrane, G-protein coupled receptor (GPCR) family. It is expressed on the surface of pituitary gonadotrope cells as well as lymphocytes, breast, ovary, and prostate. Following binding of gonadotropin-releasing hormone, the receptor associates with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Activation of the receptor ultimately causes the release of gonadotropic luteinizing hormone

(LH) and follicle stimulating hormone (FSH). Defects in this gene are a cause of

hypogonadotropic hypogonadism (HH). Alternative splicing results in multiple transcript variants encoding different isoforms. More than 18 transcription initiation sites in the 5' region and multiple polyA signals in the 3' region have been identified for this gene. [provided

by RefSeq, Jul 2008]

