

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP724109

Human CCR3 Protein, hFc Tag

Product data:

| Product Type: | Recombinant Proteins |
|------------------------|---|
| Description: | Human CCR3 Protein, hFc Tag |
| Expression Host: | HEK293 |
| Tag: | C-Human Fc |
| Predicted MW: | The protein has a predicted molecular mass of 29.9 kDa after removal of the signal peptide. The apparent molecular mass of CCR3-hFc is approximately 35-55 kDa due to glycosylation. |
| Purity: | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Reconstitution Method: | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. |
| Storage: | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Stability: | 12 months from date of despatch |
| Synonyms: | C C CKR3, C-C CKR-3, CC-CKR-3, CCR-3, CKR 3, Eosinophil eotaxin receptor, CD193, CKR3 |
| Summary: | The protein encoded by this gene is a receptor for C-C type chemokines. It belongs to family 1 of the G protein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other chemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alternatively spliced transcript variants have been described. [provided by RefSeq, Sep 2009] |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US