

Product datasheet for TP724113

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Human ZP3 Protein, hFc Tag

Product data:

Product Type: Recombinant Proteins

Description: Human ZP3 Protein, hFc Tag

Expression Host: HEK293

Tag: C-Human Fc

Predicted MW: The protein has a predicted molecular mass of 66.6 kDa after removal of the signal peptide.

The apparent molecular mass of ZP3-hFc is approximately 70-100 kDa due to glycosylation.

Purity: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie

blue staining.

Reconstitution Method: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Stability: 12 months from date of despatch
Synonyms: Sperm receptor, ZP3A/ZP3B, Zp-3

Summary: The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is

composed primarily of three or four glycoproteins with various functions during fertilization

and preimplantation development. The protein encoded by this gene is a structural component of the zona pellucida and functions in primary binding and induction of the sperm acrosome reaction. The nascent protein contains a N-terminal signal peptide sequence, a conserved ZP domain, a C-terminal consensus furin cleavage site, and a transmembrane domain. It is hypothesized that furin cleavage results in release of the mature protein from the plasma membrane for subsequent incorporation into the zona pellucida matrix. However, the requirement for furin cleavage in this process remains controversial based on mouse studies. A variation in the last exon of this gene has previously

served as the basis for an additional ZP3 locus; however, sequence and literature review reveals that there is only one full-length ZP3 locus in the human genome. Another locus encoding a bipartite transcript designated POMZP3 contains a duplication of the last four exons of ZP3, including the above described variation, and maps closely to this gene.

[provided by RefSeq, Jul 2008]

