

## **Product datasheet for TP724132**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Human CD160 Protein, hFc Tag**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Human CD160 Protein, hFc Tag

**Expression Host:** HEK293

Tag: C-Human Fc

**Predicted MW:** The protein has a predicted molecular mass of 40.9 kDa after removal of the signal

peptide. The apparent molecular mass of CD160-hFc is approximately 55-70 kDa due to

glycosylation.

**Purity:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie

blue staining.

**Reconstitution Method:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

**Stability:** 12 months from date of despatch

Synonyms: BY55, NK1, NK28

**Summary:** CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody

BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T

lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single lg-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells, spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16 cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab CD8brightCD95 CD56 CD28-CD27-cells. In tissues, CD160 is expressed on all intestinal

intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and

nonclassical MHC class I molecules. [provided by RefSeq, Jul 2008]

