

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TP724153

## Human PSMA Protein, hFc Tag

## **Product data:**

| Product Type:          | Recombinant Proteins  |
|------------------------|---|
| Description:           | Human PSMA Protein, hFc Tag   |
| Expression Host:       | HEK293  |
| Tag:                   | N-Human Fc  |
| Predicted MW:          | The protein has a predicted molecular mass of 105.7 kDa after removal of the signal peptide.The apparent molecular mass of hFc-PSMA is approximately 130 kDa due to glycosylation.  |
| Purity:                | The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie<br>blue staining.   |
| Reconstitution Method: | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants<br>before lyophilization.   |
| Storage:               | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| Stability:             | 12 months from date of despatch   |
| Synonyms:              | FGCP, FOLH, GCP2, GCPII, mGCP, NAALAD1, NAALAdase, PSM, PSMA  |
| Summary:               | This gene encodes a type II transmembrane glycoprotein belonging to the M28 peptidase<br>family. The protein acts as a glutamate carboxypeptidase on different alternative substrates,<br>including the nutrient folate and the neuropeptide N-acetyl-I-aspartyl-I-glutamate and is<br>expressed in a number of tissues such as prostate, central and peripheral nervous system<br>and kidney. A mutation in this gene may be associated with impaired intestinal absorption of<br>dietary folates, resulting in low blood folate levels and consequent hyperhomocysteinemia.<br>Expression of this protein in the brain may be involved in a number of pathological<br>conditions associated with glutamate excitotoxicity. In the prostate the protein is up-<br>regulated in cancerous cells and is used as an effective diagnostic and prognostic indicator of<br>prostate cancer. This gene likely arose from a duplication event of a nearby chromosomal<br>region. Alternative splicing gives rise to multiple transcript variants encoding several different<br>isoforms. |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US