

Product datasheet for TP724168

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Human TLR3 Protein, mFc Tag

Product data:

Product Type: Recombinant Proteins

Description: Human TLR3 Protein, mFc Tag

Expression Host: HEK293

Tag: C-Mouse Fc

Predicted MW: The protein has a predicted molecular mass of 103.5 kDa after removal of the signal peptide.

The apparent molecular mass of TLR3-mFc is approximately 130-180 kDa due to

glycosylation.

Purity: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie

blue staining.

Reconstitution Method: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Stability: 12 months from date of despatch

Synonyms: TLR3

Summary: The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which

plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta and pancreas, and is restricted to the dendritic subpopulation of the leukocytes. It recognizes dsRNA associated with viral infection, and induces the activation of NF-kappaB and the production of type I interferons. It may thus play a role in host defense against viruses. Use of alternative polyadenylation sites to generate different length transcripts has been noted for this gene. [provided by RefSeq, Jul 2008]