

## **Product datasheet for TP724219**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Human TNFRSF25 Protein, hFc Tag**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Human TNFRSF25 Protein, hFc Tag

**Expression Host:** HEK293

Tag: C-Human Fc

**Predicted MW:** The protein has a predicted molecular mass of 45.0 kDa after removal of the signal peptide.

The apparent molecular mass of TNFRSF25-hFc is approximately 55-70 kDa due to

glycosylation.

**Purity:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie

blue staining.

**Reconstitution Method:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

**Stability:** 12 months from date of despatch

**Summary:** The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor

is expressed preferentially in the tissues enriched in lymphocytes, and it may play a role in regulating lymphocyte homeostasis. This receptor has been shown to stimulate NF-kappa B activity and regulate cell apoptosis. The signal transduction of this receptor is mediated by various death domain containing adaptor proteins. Knockout studies in mice suggested the role of this gene in the removal of self-reactive T cells in the thymus. Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported, most of which are potentially secreted molecules. The alternative splicing of this gene in B and T cells encounters a programmed change upon T-cell activation, which predominantly produces

full-length, membrane bound isoforms, and is thought to be involved in controlling lymphocyte proliferation induced by T-cell activation. [provided by RefSeq, Jul 2008]