

## **Product datasheet for TP724275**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Human NPC1L1 Protein, His Tag**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Human NPC1L1 Protein, His Tag

Expression Host: HEK293
Tag: C-6×His

**Predicted MW:** The protein has a predicted molecular mass of 28.8 kDa after removal of the signal peptide.

The apparent molecular mass of NPC1L1-His is approximately 35-55 kDa due to glycosylation.

**Purity:** The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie

blue staining.

**Reconstitution Method:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

**Stability:** 12 months from date of despatch

**Summary:** The protein encoded by this gene is a multi-pass membrane protein. It contains a conserved

N-terminal Niemann-Pick C1 (NPC1) domain and a putative sterol-sensing domain (SSD) which includes a YQRL motif functioning as a plasma membrane to trans-Golgi network transport signal in other proteins. This protein takes up free cholesterol into cells through vesicular endocytosis and plays a critical role in the absorption of intestinal cholesterol. It also has the ability to transport alpha-tocopherol (vitamin E). The drug ezetimibe targets this protein and inhibits the absorption of intestinal cholesterol and alpha-tocopherol. In addition, this protein may play a critical role in regulating lipid metabolism. Polymorphic

variations in this gene are associated with plasma total cholesterol and low-density lipoprotein cholesterol (LDL-C) levels and coronary heart disease (CHD) risk. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Oct 2009]

